

Diffuse Reflections in Simple Polygons

Gill Barequet* Sarah M. Cannon† Eli Fox-Epstein‡ Benjamin Hescott‡
 Diane L. Souvaine‡ Csaba D. Tóth‡§¶ Andrew Winslow‡

Abstract

We prove a conjecture of Aanjaneya, Bishnu, and Pal that the maximum number of *diffuse reflections* needed for a point light source to illuminate the interior of a simple polygon with n walls is $\lfloor n/2 \rfloor - 1$. Light reflecting diffusely leaves a surface in all directions, rather than at an identical angle as with specular reflections.

1 Introduction

For a light source placed in a polygonal room with mirror walls, light rays that reach a wall at angle θ (with respect to the normal of the wall's surface) also leave at angle θ . In other words, for these *specular reflections* the angle of incidence equals the angle of reflection (see Fig. 1).




Figure 1: Two types of reflections. Specular reflection occurs on mirrored surfaces (left) and diffuse reflection occurs on matte surfaces (right).

Klee asked whether the interior of any room defined by a simple polygon with mirrored walls is completely illuminated by placing a single point light anywhere in the interior [7]. Tokarsky [9] gave a negative answer to this question by constructing simple polygons and pairs of points (s, t) , such that there is no path from s to t with specular reflections off the walls of the room.

On the other hand, if the walls of the polygonal room P reflect light diffusely in all directions, then it is easy to see that every point in P is illuminated after at most n *diffuse reflections* (Fig. 2). For diffuse reflections, we assume that the vertices of P absorb light, and that light does not propagate along the edges of P . A *diffuse reflection path* is a polygonal path γ contained in P

*Department of Computer Science, Technion, Haifa, Israel, barequet@cs.technion.ac.il

†Mathematical Institute, University of Oxford, Oxford, United Kingdom, cannon@maths.ox.ac.uk

‡Department of Computer Science, Tufts University, Medford, MA, {ef,hescott,dls,awinslow}@cs.tufts.edu

§California State University, Northridge, CA, USA

¶University of Calgary, Calgary, AB, Canada, cdtoth@ucalgary.ca




Figure 2: The regions of the polygon illuminated by a light source s after 0, 1, 2, and 3 diffuse reflections.

such that every interior vertex of γ lies in the relative interior of some edge of P , and the relative interior of every edge of γ is in the interior of P .

Denote by $V_k(s) \subseteq P$ the part of the polygon illuminated by a light source s after at most k diffuse reflections. Formally, $V_k(s)$ is the set of points $t \in P$ such that there is a diffuse reflection path from s to t with at most k interior vertices. Hence, $V_0(s)$ is the visibility region of point s within the polygon P and so is a simply connected region with $O(n)$ edges. Aronov et al. [3] showed that $V_1(s)$ is simply connected with at most $O(n^2)$ edges, and this bound cannot be improved. Brahma et al. [5] constructed simple polygons and a source s such that $V_2(s)$ is not simply connected, and showed that $V_3(s)$ can have as many as $\Omega(n)$ holes. Extending the work of [3], Aronov et al. [2, 4] and Prasad et al. [8] bounded the complexity of $V_k(s)$ at $O(n^9)$ and $\Omega(n^2)$ for all k . It remains an open problem to close the gap between these bounds for $k \geq 2$.

Finding a shortest diffuse illumination path between two given points in a simple polygon by brute force is possible in $O(n^{10})$ time using the result of Aronov et al. [4]. Ghosh et al. [6] presented a 3-approximation in a much-improved $O(n^2)$ time, and their approximation applies even if the polygon P has holes.




Figure 3: Left: An orthogonal spiral polygon with $n = 20$ vertices [1], where every diffuse reflection path between s and t has at least $\lceil n/2 \rceil - 2 = 8$ turns. Right: A zig-zag polygon with $n = 16$ vertices where every diffuse reflection path between s and t has at least $\lfloor n/2 \rfloor - 1 = 7$ reflections.

Results. We determine the maximum length of a diffuse illumination path in a simple polygon in n vertices. Let $D(n)$ be the smallest integer $k \in \mathbb{N}_0$ such that for any simple polygon P with n vertices and any two interior points $s, t \in \text{int}(P)$, there is a diffuse illumination path between s and t with at most k interior vertices (i.e., at most k reflections). Aanjaneya et al. [1] conjectured that $D(n) \leq \lceil n/2 \rceil - 1$ and construct an example which yields $D(n) \geq \lfloor n/2 \rfloor - 2$; see Fig. 3 (left). The zig-zag polygon in Figure 3 (right) shows that $D(n) \geq \lfloor n/2 \rfloor - 1$. Here we prove that $D(n) = \lfloor n/2 \rfloor - 1$.

Theorem 1. *Let P be a simple polygon with $n \geq 3$ vertices. We have $\text{int}(P) \subseteq V_k(s)$ for every $s \in \text{int}(P)$ and $k \geq \lfloor n/2 \rfloor - 1$.*

Corollary 1. *Let P be a simple polygon with $n \geq 3$ vertices. Between any two points $s, t \in \text{int}(P)$, there exists a diffuse reflection path with at most $\lfloor n/2 \rfloor - 1$ reflections.*

Definitions. For a planar set $S \subseteq \mathbb{R}^2$, let $\text{int}(S)$ and $\text{cl}(S)$ denote the set of interior points of S and the closure of S , respectively. The boundary of S , denoted ∂S , is $\partial S = \text{cl}(S) \setminus \text{int}(S)$. Let P be a simple polygon with $n \geq 3$ vertices. A *chord* of P is a line segment ab , such that $a, b \in \partial P$ and the relative interior of ab lies in $\text{int}(P)$. The visibility polygon of a chord ab , $V_0(ab)$, is the set of points visible from a point in ab (i.e., the *weak visibility polygon* of ab). A set $S \subseteq P$ *weakly covers* an edge e of P if S intersects the relative interior of e .

2 A Set of Regions R_k

Let P be a simple polygon with n vertices, and let $s \in \text{int}(P)$. Let us assume that the vertices of P and s are in general position, that is, there are only trivial algebraic relations among s and the vertices of P . (This assumption simplifies the presentation, but it is not essential for the proof.)

Instead of tackling $V_k(s)$ directly, we define an infinite sequence of simply-connected regions $R_0 \subseteq R_1 \subseteq R_2 \subseteq \dots$, such that $R_0 = V_0(s)$ and $R_k \subseteq V_k(s)$ for all $k \in \mathbb{N}$, and then show that $\text{int}(P) \subseteq R_{\lfloor n/2 \rfloor - 1}$. For every $k \in \mathbb{N}_0$, the region R_k will have the following structural properties.

- (i) The boundary of R_k is a simple polygon, in which each edge is either a chord of P (called a *window*) or part of an edge of P . (The boundary of R_k is not necessarily part of R_k .)
- (ii) The windows of R_k are pairwise disjoint.
- (iii) For each window ab , one endpoint, say a , is a reflex vertex of P , and the other endpoint b lies in the relative interior of some edge e_{ab} of P .
- (iv) For each window ab , we have $b \notin R_k$, but in any neighborhood of b , some point on the edge e_{ab} is in R_k .

It is easy to verify that all these properties hold for the initial region $R_0 = V_0(s)$ (e.g., see Figure 4, left). Every window ab of P decomposes P into two simple polygons sharing the side ab . Denote by U_{ab} the polygon that does not contain R_k . A window ab of R_k is *saturated* if one endpoint of every chord of P crossing ab is in R_k ; otherwise, it is *unsaturated*. Recall that for a point $s \in P$, we denote by $V_0(s)$ the set of points in P visible from s . Note that if s lies in the interior of an edge e of P , then $V_0(s) \cap e = \{s\}$, that is, s is the only point in e visible from s .

Construction of the regions R_k . We are now ready to define the sets R_k , $k \in \mathbb{N}_0$. Refer to Fig. 4. Let $R_0 = V_0(s)$. If $\text{int}(P) \subseteq R_k$, then let $R_{k+1} = \text{cl}(R_k) = P$. If $\text{int}(P) \not\subseteq R_k$, then R_k has at least one window. For each window ab , we define a set W_{ab} as follows. If ab is saturated, then let $W_{ab} = V_0(ab) \cap U_{ab}$. If ab is unsaturated, then let $c \in R_k \cap \partial P$ be a point close to b on e_{ab} such that no line determined by two vertices of P separates b and c ; and then let $W_{ab} = V_0(c) \cap U_{ab}$. Let R_{k+1} be the union of $\text{cl}(R_k)$ and the sets W_{ab} for all windows ab of R_k .

Proposition 1. *If $R_k \subseteq V_k(s)$, then $\text{cl}(R_k) \subseteq V_{k+1}(s)$ for all $k \in \mathbb{N}_0$*




Figure 4: Left: A simple polygon P with illumination point s . The visibility polygon $V_0(s)$ has three windows a_1b_1 , a_2b_2 , and a_3b_3 . Window a_1b_1 is saturated, while a_2b_2 and a_3b_3 are unsaturated. Right: R_1 is the union of the closure of $R_0 = V_0(s)$ and visibility polygons of each of its three windows.

Proof. Let $p \in \partial R_k \setminus R_k$ be a point on the boundary of R_k that is not in R_k . Since $p \in \text{cl}(R_k) \subseteq \text{cl}(V_k(s))$, we have $s \in \text{cl}(V_k(p))$ by symmetry. Let $s_\varepsilon \in V_k(p)$ be a point in a small ε -neighborhood of s such that there is a diffuse reflection path $(p, r_\varepsilon(1), \dots, r_\varepsilon(\ell), s_\varepsilon)$ with $\ell \leq k$, where the points $r_\varepsilon(1), \dots, r_\varepsilon(\ell)$ lie in the interior of some edges of P . We may assume that $s_\varepsilon \rightarrow s$ and $r_\varepsilon(\ell) \rightarrow r$ as $\varepsilon \rightarrow 0$. Let q be a point where ray $\overrightarrow{rs}$ hits the boundary of P (recall that $s \in \text{int}(P)$). The ray $\overrightarrow{r_\varepsilon(\ell)s_\varepsilon}$ hits ∂P in a small neighborhood of q if $\varepsilon > 0$ is sufficiently small. Hence in every δ -neighborhood of q , there is a point $r_\delta \in \partial P$ that lies in both $V_k(p)$ and the interior of an edge of P . If $\delta > 0$ is sufficiently small, then r_δ directly sees s , and so there is a diffuse reflection path of length at most $k + 1$ between p and s . $\square$

The next two propositions count the new edges of P weakly covered by R_{k+1} in U_{ab} for a window ab of R_k .

Proposition 2. *Let ab be a saturated window of R_k . Then $W_{ab} = V_0(ab) \cap U_{ab}$ weakly covers at least one edge of P not weakly covered by R_k . Furthermore, if $U_{ab} \not\subseteq V_0(ab)$, then $V_0(ab) \cap U_{ab}$ weakly covers at least two edges of P not weakly covered by R_k .*

Proof. By property (iii), a is a reflex vertex of P . Denote by ad the edge of P adjacent to U_{ab} . It is clear that R_k does not weakly cover ad , but $V_0(ab)$ weakly covers ad . This proves the first claim.

By property (iv), b lies on an edge e_{ab} of P . Now consider all the chords of P that cross ab and are parallel to ad or to e_{ab} . If all chords parallel to ad have an endpoint in e_{ab} and all chords parallel to e_{ab} have an endpoint on ab , then U_{ab} is the triangle $\Delta(abd)$, and $U_{ab} \subseteq V_0(ab)$. Otherwise, one of the chords has an endpoint on some edge of P that is disjoint from R_k . In this case $V_0(ab) \cap U_{ab}$ weakly covers at least two edges of P not weakly covered by R_k . $\square$

Proposition 3. *Let ab be an unsaturated window of R_k , and let $c \in R_k \cap \partial P$ be a point so close to b on the boundary of P such that no line determined by two vertices of P separates b and c . Then $V_0(c) \cap U_{ab}$ weakly covers at least one edge of P not weakly covered by R_k .*

Proof. By property (iii), a is a reflex vertex of P . Denote by ad the edge of P adjacent to U_{ab} . It is clear that R_k does not weakly cover ad , but $V_0(c)$ weakly covers ad . $\square$

Lemma 1. $R_k \subseteq V_k(s)$ for all $k \in \mathbb{N}_0$.

Proof. We show by induction on $k \geq 0$ that $R_k \subseteq V_k(s)$ for all $k \in \mathbb{N}_0$. In the base case, we have $R_0 = V_0(s)$ by definition. The Propositions 1, 2, and 3 together imply that if $R_k \subseteq V_k(s)$, then $R_{k+1} \subseteq V_{k+1}(s)$ for all $k \in \mathbb{N}_0$. $\square$

3 Counting Weakly Covered Edges in R_k

The proof of Theorem 1 is based on counting the edges in the polygon weakly covered by R_k , which we denote μ_k . It is not difficult to show that $R_0 = V_0$ weakly covers at least 3 edges ($\mu_0 \geq 3$). We show that the invariant

$$\mu_k \geq \min(2k + 3, n). \quad (1)$$

is maintained for all $k \in \mathbb{N}_0$, which immediately implies Theorem 1. Invariant (1) is clearly maintained when the number of edges weakly covered by R_k increases by two or more. Unfortunately, this is not always the case: in some instances, R_{k+1} weakly covers only one more edge than R_k (i.e., $\mu_{k+1} = \mu_k + 1$). We introduce the notion of “critical” cases when $\mu_k = 2k + 3$ and by careful analysis show that for every critical R_k , R_{k+1} weakly covers at least two new edges of P , maintaining invariant (1).

Let λ_k denote the number of windows of R_k . Since the regions R_k monotonically increase (i.e., $R_{k-1} \subseteq R_k$, $k \in \mathbb{N}_0$), we have $\mu_k \leq \mu_{k+1}$ for all $k \in \mathbb{N}_0$. By Propositions 2 and 3, we have

$$\mu_{k+1} \geq \mu_k + \lambda_k \quad \text{for all } k \in \mathbb{N}_0. \quad (2)$$

In particular, this implies $\mu_k < \mu_{k+1}$ until $\mu_k = n$. The initial value μ_0 is at least three.

Proposition 4. *If $s \in \text{int}(P)$, then $V_0(s)$ weakly covers at least three edges of P .*

Proof. For every edge e of P , let $H(e)$ be a halfplane bounded by the line through e and containing s . Consider the intersection C of the halfplanes $H(e)$ for all edges e weakly visible from s . Then C is a convex region containing s . Note that C is bounded, otherwise it would contain an unbounded ray emitted by s which would hit the relative interior of another edge of P . Since C is a convex polygon, it has at least 3 edges, and so at least 3 edges of P are weakly visible from s . $\square$

Invariant (1) holds for $k = 0$ by Proposition 4. After some preparation we show (Lemma 3) that invariant (1) is maintained for all $k \in \mathbb{N}$.

A region R_k is called *critical* if $\mu_k = 2k + 3$ and $\mu_k < n$. From (2), it is enough to show that if R_k is critical, then R_{k+1} satisfies (1). Invariant (1) is maintained when $\mu_{k+1} = \mu_k + 2$. Invariant (2) implies that $\mu_{k+1} \geq \mu_k + 1$ while $\mu_k < n$, since if $\mu_k < n$ then $R_k \neq \text{int}(P)$. Hence, invariant (1) fails to hold for R_{k+1} only if R_k is critical and $\mu_{k+1} = \mu_k + 1$. For every critical region R_k , we establish one of the following two conditions:

- (A) All windows of R_k are saturated;
- (B) $\lambda_k \geq 2$, but R_k has an unsaturated window.

We first show that the property (A) or (B) hold for R_0 if R_0 is critical.

Proposition 5. *If R_0 is critical, then it satisfies (A) or (B).*

Proof. Suppose that $\mu_0 = 3$. If R_0 has at least two windows and $\lambda_0 \geq 2$, then condition (A) or (B) is clearly satisfied. Suppose that $\lambda_0 = 1$. Then the boundary of $R_0 = V_0(s)$ consists of a unique window ab , and segments along the boundary of P , which are directly visible from s . Hence any chord of P that crosses ab has an endpoint in $R_0 \cap \partial P$, and condition (a) is satisfied. $\square$

Proposition 6. *Suppose that R_k is noncritical but R_{k+1} is critical. Then $\lambda_k = \lambda_{k+1} = 1$, the window of R_k is unsaturated, and the window of R_{k+1} is saturated.*

Proof. Because R_k is noncritical and R_{k+1} is critical, we have $\mu_k \geq 2k+4$ and $\mu_{k+1} = 2(k+1)+3 = 2k+5$. Combined with (2), we have $\lambda_k = 1$, that is, R_k has a unique window. This unique window cannot be saturated by Proposition 2 and the fact that $\mu_{k+1} = \mu_k + 1$.




Figure 5: The situation in Proposition 6. The region R_k is noncritical and R_{k+1} is critical. The unique window ab of R_k is unsaturated, while the unique window $a'b'$ of R_{k+1} is saturated.

Refer to Figure 5. Because the window of R_k is unsaturated, $R_{k+1} = \text{cl}(R_k) \cup W_{ab}$, where $W_{ab} = V_0(c) \cap U_{ab}$ for a point $c \in e_{ab}$. Since $\mu_{k+1} - \mu_k = 1$, region R_{k+1} weakly covers only one additional edge. So the supporting line of edge e_{ab} intersects edge ad at some point b' .

Denote by a' the endpoint of e_{ab} that lies in segment bb' . Observe that $V_{ab} = \Delta(acb') \setminus cb'$, and so R_{k+1} has a unique window $a'b'$. Every chord of P that crosses $a'b'$ has an endpoint in either the relative interior of ab' or in $\text{cl}(R_k)$, which are contained in R_{k+1} , and so $a'b'$ is saturated. $\square$

Proposition 7. *If $\lambda_k = \lambda_{k+1} = 1$ and the unique window of R_k is saturated, then the unique window of R_{k+1} is also saturated.*

Proof. Let ab be the saturated window of R_k , and let $a'b'$ be a window of R_{k+1} . By construction $W_{ab} = V_0(ab) \cap U_{ab}$, and $\text{cl}(R_k) \cup W_{ab} \subseteq R_{k+1}$. Note that $a'b'$ lies in U_{ab} . Consider a chord cd of P that crosses $a'b'$ with $d \in U_{a'b'}$. We need to show that $c \in R_{k+1}$. If cd crosses ab , then $c \in R_k \subseteq R_{k+1}$ since ab is saturated. If $c = a$ or $c = b$ then $c \in R_{k+1}$ since $a, b \in \text{cl}(R_k) \subseteq R_{k+1}$. Otherwise $c \in \partial W_{ab} \setminus (ab \cup a'b')$. Since ab is saturated, all points of $\partial W_{ab} \setminus (ab \cup a'b')$ are in $W_{ab} \subseteq R_{k+1}$. In all cases, $c \in R_{k+1}$, and so the window $a'b'$ is saturated. $\square$

Proposition 8. *If both R_0 and R_1 are critical and $\lambda_0 = 2$, then all windows of R_1 are saturated.*

Proof. If both R_0 and R_1 are critical, then $\mu_0 = 3$ and $\mu_1 = 5$. Denote by a_1b_1 and a_2b_2 the two windows of R_0 . Neither window of R_0 is saturated, otherwise we would have $\mu_1 > 5$ by (2) and Propositions 2 and 3. The boundary of R_0 consists of 5 line segments: the windows a_1b_1 , a_2b_2 and three segments along three edges of P weakly covered by R_0 .

Property (ii) implies that a_1b_1 and a_2b_2 are not adjacent, hence they are both incident to some edge e of P weakly covered by R_0 . Property (iii) implies that a_1 and a_2 are reflex vertices of P . Since

$R_0 = V_0(s)$ is the visibility region of s , both a_1 and a_2 are reflex vertices in R_0 . If a_1 or a_2 is incident to e , then both windows are saturated, contradicting our earlier observation. Therefore, neither a_1 nor a_2 is incident to e , hence both b_1 and b_2 lie on e . Then $R_1 = \text{cl}(R_0) \cup V_0(c_1) \cup V_0(c_2)$, where $c_1, c_2 \in e$. Since R_1 weakly covers exactly one new edge of P behind each of window, every window of R_1 is collinear with edge e . It follows that a chord of P that crosses any window of R_1 cannot have an endpoint on edge e . Therefore one endpoint of such a chord is in $R_1 = \text{cl}(R_0) \cup V_0(c_1) \cup V_0(c_2)$, and all windows of R_1 are saturated. $\square$

Proposition 9. *Suppose that R_k , R_{k+1} and R_{k+2} are critical, $\lambda_k = 1$ and $\lambda_{k+1} = 2$. Then all windows of R_{k+2} are saturated.*

Proof. Let ab be the unique window of R_k and let $a'_1b'_1$ and $a'_2b'_2$ be the two windows of R_{k+1} . By property (iii), a is a reflex vertex of P . Denote by ad the edge of P adjacent to U_{ab} and a . By property (iv), b lies on an edge e_{ab} of P . Note that both $a'_1b'_1$ and $a'_2b'_2$ are on the boundary of W_{ab} . Since R_k and R_{k+1} are both critical, we have $\mu_{k+1} = \mu_k + 2$. So R_{k+1} weakly covers two new edges of P : one is ad and call the other e . Now the boundary of W_{ab} is formed by segments ab , $a'_1b'_1$, $a'_2b'_2$, and parts of the edges ad , e_{ab} and e . Property (ii) implies that $a'_1b'_1$ and $a'_2b'_2$ are disjoint, and by construction they are both disjoint from ab as well. Without loss of generality, we may assume that ∂W_{ab} contains, in counterclockwise order, parts of ab , e_{ab} , $a'_1b'_1$, e , $a'_2b'_2$, and ad .




Figure 6: R_k has a unique window ab , and V_{ab} weakly covers two new edges of P : ad and e . R_{k+1} has two windows $a'_1b'_1$ and $a'_2b'_2$. Left: edge e is incident to b'_1 and a'_2 ; no chord of P crosses both $a'_1b'_1$ and $a'_2b'_2$ and so both are saturated. Right: edge e is incident to b'_1 and b'_2 . The two windows of R_{k+2} are collinear with edge e and so both are saturated.

Since both R_{k+1} and R_{k+2} are critical, we have $\mu_{k+2} = \mu_{k+1} + 2$. That is, R_{k+2} weakly covers exactly one new edge of P behind each of the two windows $a'_1b'_1$ and $a'_2b'_2$, namely the edges of P incident to a'_1 and a'_2 , respectively. It follows that R_{k+2} has at most two windows: at most one behind each of $a'_1b'_1$ and $a'_2b'_2$.

Property (iii) implies that a'_1 and a'_2 are reflex vertices of P . Since W_{ab} is the visibility region of a point or a segment, both a'_1 and a'_2 are reflex vertices in W_{ab} , and hence in R_{k+1} . We distinguish two cases depending on the number of points in $\{a'_1, a'_2\}$ incident to edge e . Refer to Fig. 6.

Case 1: a'_1 or a'_2 is incident to e . Then no chord of P intersects both $a'_1b'_1$ and $a'_2b'_2$. In this case, both $a'_1b'_1$ and $a'_2b'_2$ are saturated. Hence any window of R_{k+2} (at most one behind each of $a'_1b'_1$ and $a'_2b'_2$) is also saturated.

Case 2: neither a'_1 nor a'_2 is incident to e . Then both b'_1 and b'_2 lie on edge e . In this case neither $a'_1b'_1$ nor $a'_2b'_2$ is saturated. Then $R_{k+2} = \text{cl}(R_{k+1}) \cup V_0(c'_1) \cup V_0(c'_2)$, where $c'_1, c'_2 \in e$ are points close to b'_1 and b'_2 , respectively. Since R_{k+2} weakly covers exactly one new edge of P behind each of $a'_1b'_1$ and $a'_2b'_2$, every window of R_{k+2} is collinear with edge e . It follows that a chord of P that crosses any window of R_{k+2} cannot have an endpoint on edge e . Therefore one endpoint of such a chord is in $R_{k+2} = \text{cl}(R_{k+1}) \cup V_0(c'_1) \cup V_0(c'_2)$. Hence any window of R_{k+2} is saturated. $\square$

Lemma 2. *Let $R_h, \dots, R_k$ be a maximal sequence of critical regions. Then, condition (A) or (B) applies for each $i = h, \dots, k$.*

Proof. We have either $h = 0$ and then R_0 satisfies (A) or (B) by Proposition 5; or $h \geq 1$ and R_h satisfies (A) by Proposition 6. We now show successively for $i = h + 1, \dots, k$ that R_i satisfies (A) or (B).

Suppose that both R_i and R_{i+1} are critical and R_i satisfies (A). We have $\mu_{i+1} = \mu_i + 2$ since both R_i and R_{i+1} are critical. By Proposition 2, $\mu_{i+1} \geq \mu_i + \lambda_i + 1$ and so $\lambda_i = 1$. If $\lambda_{i+1} = 1$, then R_{i+1} satisfies (A) by Proposition 7. If $\lambda_{i+1} \geq 2$, then R_{i+1} satisfies (B).

If both R_0 and R_1 are critical and R_0 satisfies (B), then R_1 satisfies (A) by Proposition 8. Suppose that R_i, R_{i+1} , and R_{i+2} are critical, R_i satisfies (A), and R_{i+1} satisfies (B). Note $\lambda_{i+1} = 2$, otherwise R_{i+2} is not critical. Then R_{i+2} satisfies (A) by Proposition 9. It follows that no two consecutive critical regions satisfy (B); and whenever R_{i+1} satisfies (B), then R_{i+2} satisfies (A). $\square$

Lemma 3. *Invariant (1) holds for all $k \in \mathbb{N}_0$.*

Proof. It is enough to show that whenever R_k is critical, the region R_{k+1} satisfies (1). Consider a maximal sequence $R_h, \dots, R_k$ of critical regions such that $\mu_{k+1} < n$. By Lemma 2, we have $\lambda_k \geq 2$ or all windows of R_k are saturated. If $\lambda_k \geq 2$, then $\mu_{k+1} \geq \mu_k + 2$ by (2). If all windows of R_k are saturated, then R_{k+1} weakly covers at least one new edge of P behind each window of R_k , and at least two edges behind one of the windows due to Proposition 2 and the condition $\mu_{k+1} < n$. Therefore, $\mu_{k+1} \geq \mu_k + \lambda_k + 1$. In both cases, we have $\mu_{k+1} \geq \mu_k + 2 \geq (2k + 3) + 2 = 2(k + 1) + 3$, and R_{k+1} satisfies (1), as required. $\square$

Theorem 1 Let P be a simple polygon with $n \geq 3$ vertices. We have $\text{int}(P) \subseteq V_k(s)$ for every $s \in \text{int}(P)$ and $k \geq \lfloor n/2 \rfloor - 1$.

Proof. If $R_{\lfloor n/2 \rfloor - 1}$ has a window ab , then by property (iii) there is an edge ad not weakly covered by $R_{\lfloor n/2 \rfloor - 1}$, contradicting invariant (1). Therefore $R_{\lfloor n/2 \rfloor - 1}$ has no windows, and $\text{int}(P) \subseteq R_{\lfloor n/2 \rfloor - 1}$, as claimed. $\square$

Acknowledgements

We would like to thank the LAGOS reviewers for helpful comments improving the presentation of the paper and the MIT-Tufts Research Group on Computational Geometry for initial discussions of the problem.

References

- [1] M. Aanjaneya, A. Bishnu, and S. P. Pal, Directly visible pairs and illumination by reflections in orthogonal polygons, *Proc. 24th European Workshop on Computational Geometry*, Nancy, France, 241–244, March, 2008.
- [2] B. Aronov, A. Davis, T. K. Dey, S. P. Pal, and D. C. Prasad, Visibility with multiple reflections, *Discrete & Computational Geometry* **20** (1998), 61–78.
- [3] B. Aronov, A. Davis, T. K. Dey, S. P. Pal, and D. C. Prasad, Visibility with one reflection, *Discrete & Computational Geometry* **19** (1998), 553–574.
- [4] B. Aronov, A. Davis, J. Iacono, and A. S. C. Yu, The complexity of diffuse reflections in a simple polygon, *7th LATIN*, vol. 3887 of LNCS, Springer, 2006, pp. 93–104.
- [5] S. Brahma, S. P. Pal, and D. Sarkar, A linear worst-case lower bound on the number of holes inside regions visible due to multiple diffuse reflections, *J. of Geometry* **81** (2004), 5–14.
- [6] S. K. Ghosh, P. P. Goswami, A. Maheshwari, S. C. Nandy, S. P. Pal, and S. Sarvattomananda, Algorithms for computing diffuse reflection paths in polygons, *Vis. Comput.* **28** (2012), 1229–1237.
- [7] V. Klee, Some unsolved problems in plane geometry, *Mathematics Magazine* **52**(3) (1979), 131–145.
- [8] D. C. Prasad, S. P. Pal, and T. K. Dey, Visibility with multiple diffuse reflections. *Computational Geometry* **10**(3) (1998), 187–196.
- [9] G. Tokarsky, Polygonal rooms not illuminable from every point, *The American Mathematical Monthly* **102**(10) (1995), 867–879.